

Supplementary Table S2. Overall infection status of *Clonorchis sinensis* metacercariae in fish in all 3 locations from Banbyeon-cheon, Yeongyang-gun, Gyeongsangbuk-do, Korea

Fish species infected	No. of fish examined	Infection rate (%)	No. of DTM detected	
			Total	Average (range)
Hyeong-ri (site A)				
<i>Pungtungia herzi</i>	34	9 (26.5)	149	16.6 (1–109)
<i>Sarcocheilichthys nigripinnis morii</i>	3	1 (33.3)	8	8.0
<i>Zacco platypus</i>	34	2 (5.9)	2	1.0
Subtotal	71	12 (16.9)	159	13.3
Hapgang-ri (site B)				
<i>Pungtungia herzi</i>	16	5 (31.3)	301	60.2 (3–26)
<i>Acheilognathus majusculus</i>	1+11	2 (18.2)	11	5.5 (1–4)
<i>Acheilognathus lanceolatus</i>	21+9	1 (11.1)	8	8
<i>Sarcocheilichthys nigripinnis morii</i>	5	4 (80.0)	17	4.25(1–4)
<i>Pseudogobio esocinus</i>	2	1 (50)	6	6
<i>Squalidus japonicus koreanus</i>	4	1 (25)	26	26
Subtotal	47	14 (29.8)	369	26.4
Total	118	26 (22.0)	528	20.3 (1–109)
Bangjeon-ri (site C)				
<i>Pungtungia herzi</i>				
<i>Squalidus japonicus koreanus</i>	1	1 (100)	1	1
<i>Coreoperca herzi</i>	1	1 (100)	1	1
Subtotal	2	2 (100)	2	1
Total	120	28 (23.3)	530	19 (1–109)