

Supplementary Table S3. Overall infection status of *Centrocestus armatus* metacercariae in fish in all 3 locations from Banbyeon-cheon, Yeongyang-gun, Gyeongsangbuk-do, Korea

Fish species infected	No. of fish examined	Infection rate (%)	No. of DTM detected	
			Total	Average (range)
Hyeong-ri (site A)				
<i>Sarcocheilichthys nigripinnis morii</i>	3	1 (33.3)	2	2.0 (1–3)
<i>Nipponocypris temminckii</i>	33	33 (100)	4,583	138.9 (4–720)
<i>Acheilognathus majusculus</i>	1	1 (100)	1	1
<i>Squalidus japonicus koreanus</i>	3	1 (33.3)	6	6
<i>Pseudogobio esocinus</i>	3	1 (33.3)	1	1
<i>Zacco platypus</i>	34	34 (100)	4,883	143.6 (3–673)
<i>Misgurnus anguillicaudatus</i>	1	1 (100)	2	2
Subtotal	78	72 (92.3)	9,478	131.6
Hapgang-ri (site B)				
<i>Pungtungia herzi</i>	16	1 (6.25)	12	
<i>Sarcocheilichthys nigripinnis morii</i>	5	2 (40)	5	2.5 (1–2)
<i>Coreoperca herzi</i>	1	2 (100)	2,135	1,068 (2–135)
<i>Zacco platypus</i>	17	14 (82.4)	234	16.7 (1–6)
<i>Squalidus japonicus koreanus</i>	4	2 (50.0)	9	4.5
Subtotal	43	21 (47.7)	2,395	114
Total	121	93 (76.8)	11,873	127 (1–720)
Bangjeon-ri (site C)				
<i>Squalidus japonicus koreanus</i>	1	1 (100)	4	4
<i>Coreoperca herzi</i>	1	1 (100)	1	1
Subtotal	2	2	5	2.5
Total	123	95 (77.2)	11,878	125 (1–720)