

Supplementary Table S7. Overall infection status of *Metacercaria hasegawai* metacercariae in fish in all locations from Banbyeon–cheon, Yeongyang-gun, Gyeongsangbuk-do, Korea

Fish species infected	No. of fish examined	Infection rate (%)	No. of DTM detected	
			Total	Average (range)
Hyeong-ri (site A)				
<i>Pungtungia herzi</i>	34	9 (26.5)	58	6.4 (1–26)
<i>Acheilognathus lanceolatus</i>	21	3 (14.3)	5	1.7 (1–3)
<i>Sarcocheilichthys nigripinnis morii</i>	3	1 (33.3)	4	4
<i>Acheilognathus majusculus</i>	1	1 (100)	6	6 (1–2)
<i>Squalidus japonicus koreanus</i>	3	2 (66.7)	5	2.5 (2–3)
<i>Zacco platypus</i>	34	1 (2.9)	1	1
<i>Coreoperca herzi</i>	5	1 (20.0)	1	1
Subtotal	101	18 (17.8)	80	4.4
Hapgang-ri (site B)				
<i>Pungtungia herzi</i>	16	9 (56.3)	175	19.4 (2–8)
<i>Acheilognathus majusculus</i>	11	11 (100)	539	49 (1–14)
<i>Acheilognathus lanceolatus</i>	9	9 (100)	1,643	182.6 (2–44)
<i>Sarcocheilichthys nigripinnis morii</i>	5	5 (100)	24	4.8 (1–2)
<i>Pseudogobio esocinus</i>	2	2 (100)	7	3.5 (1–2)
<i>Zacco platypus</i>	17	2 (11.8)	3	1.5
<i>Squalidus japonicus koreanus</i>	4	4 (100)	28	7 (1–3)
Subtotal	64	42 (65.6)	2,419	57.6
Total	165	60 (36.3)	2,499	41.7 (1–44)
Bangjeon-ri (site C)				
<i>Squalidus japonicus koreanus</i>	1	1 (100)	10	10
<i>Acheilognathus lanceolatus</i>	2	1 (50)	4	4
Subtotal	3	2 (66.7)	14	7
Total	168	62 (36.9)	2,513	41 (1–44)