

Supplementary Table S2. Infection status of DsMc in fish^a from the water systems of Han-gang

Positive fish species	No. of fish examined	No. (%) of fish infected	No. of DsMc detected	
			Range	Average
<i>Zacco platypus</i>	85	28 (32.9)	1-5	2.1
<i>Nipponocypris koreanus</i>	63	7 (11.1)	1-2	1.4
<i>Acheilognathus lanceolatus</i>	59	24 (40.7)	1-10	2.9
<i>Pseudogobio esocinus</i>	49	26 (53.1)	1-54	10.2
<i>Carassius auratus</i>	49	9 (18.4)	1-5	2.1
<i>Hemibarbus longirostris</i>	39	4 (10.3)	1-2	1.3
<i>Microphysogobio longidorsalis</i>	35	9 (25.7)	1-12	3.8
<i>Hemibarbus labeo</i>	18	14 (77.8)	1-7	3.4
<i>Coreoperca herzi</i>	18	1 (5.6)	-	1.0
<i>Cyprinus carpio</i>	2	1 (50.0)	-	4.0
<i>Opsariichthys uncirostris</i>	1	1 (100)	-	1.0
Total	418 ^b	124 (29.7)	1-54	4.2

DsMc, *Diplostomum* spp. metacercariae.

^aA total of 707 fish in 30 species from Seom-gang (37.42690; 127.89634), a branch stream of Han-gang, in Wonju-si (in 2018~2020), Gangwon-do were examined; ^bA total of 289 fish in 19 spp. were DsMc negative.