

Supplementary Table S4. Infection status of DsMc in fish^a from Mangyeong-gang in Wanju-gun, Jeollabuk-do

Positive fish species	No. of fish examined	No. (%) of fish infected	No. of DsMc detected	
			Range	Average
<i>Pungtungia herzi</i>	185	5 (2.7)	1-3	1.8
<i>Pseudogobio esocinus</i>	123	71 (57.7)	1-31	6.8
<i>Odontobutis platycephala</i>	106	8 (7.5)	1-2	1.1
<i>Zacco platypus</i>	100	44 (44.0)	1-59	10.8
<i>Hemibarbus longirostris</i>	94	15 (16.0)	1-9	1.7
<i>Coreoperca herzi</i>	76	9 (11.8)	1-3	1.6
<i>Nipponocypris temminckii</i>	61	32 (52.5)	1-32	5.7
<i>Carassius auratus</i>	32	1 (3.1)	-	1.0
<i>Nipponocypris koreanus</i>	28	20 (71.4)	2-18	6.6
<i>Acheilognathus yamatsutae</i>	13	8 (61.5)	1-9	4.0
<i>Acheilognathus lanceolatus</i>	10	8 (80.0)	1-13	5.0
<i>Squalidus chankaensis</i>	3	2 (66.7)	7-9	8.0
Total	831 ^b	223 (26.8)	1-59	6.4

DsMc, *Diplostomum* spp. metacercariae.

^aA total of 927 fish in 31 species from Soyang-cheon (36.11427; 127.58775), a branch stream of Mangyeong-gang, in Wanju-gun (462 fish of 15 spp. in 2013~2015; 465 fish of 25 spp. in 2018~2019), Jeollabuk-do were examined; ^bA total of 96 fish in 19 spp. were DsMc negative.