

Table S1. Nodules and mass forming lesions in different groups of mice

| Group(s) | Mice ID | Lobe(s) | Gray-white nodule(s) | | | | Large mass forming lesion(s) | | | |
|------------|---------|---------------|----------------------|-------------|------------|---------------------------|------------------------------|-------------|------------|---------------------------|
| | | | No. | Length (mm) | Width (mm) | Volume (mm ³) | No. | Length (mm) | Width (mm) | Volume (mm ³) |
| Ctrl. | None | | | | | | | | | |
| Cs | 2-2 | Left | 1 | 0.7 | 0.7 | 0.18 | | | | |
| | 2-4 | Right | 1 | 1.5 | 0.8 | 0.48 | | | | |
| | 2-6 | Right, middle | 1 | 3.0 | 2.7 | 10.93 | 1 | 10.00 | 7.5 | 281.25 |
| NDMA | None | | | | | | | | | |
| Cs+NDMA | None | | | | | | | | | |
| DC | None | | | | | | | | | |
| Cs+DC | 6-2 | Caudate | | | | | 1 | 10.25 | 7.4 | 280.64 |
| | 6-5 | Left | 1 | 1.3 | 1.1 | 0.79 | | | | |
| NDMA+DC | 7-7 | Right | 1 | 0.8 | 0.5 | 0.10 | | | | |
| | 7-10 | Left | 1 | 0.7 | 0.5 | 0.09 | | | | |
| Cs+NDMA+DC | 8-6 | Middle | 1 | 0.9 | 0.8 | 0.29 | | | | |
| | 8-7 | Middle | 1 | 2.2 | 2.2 | 5.32 | | | | |
| | 8-7 | Left | 1 | 0.5 | 0.5 | 0.06 | | | | |
| | 8-7 | Left | 1 | 0.5 | 0.2 | 0.02 | | | | |
| | 8-13 | Middle | 1 | 1.2 | 1.2 | 0.99 | | | | |