

**Supplementary Table S1.** Yearly infection status of *Clonorchis sinensis* metacercariae (CsMc) in fishes from the irrigation canal of To-gyo-jeosujji in Cheorwon-gun, Gangwon-do, Korea

Year and fish sp. examined	No. of fish examined	No. (%) of fish infected	No. of CsMc detected	
			Range	Average
2018				
<i>Zacco platypus</i>	54	20 (37.0)	1-22	4.3
<i>Acheilognathus rhombeus</i>	25	25 (100)	1-45	9.9
<i>Pseudorasbora parva</i>	24	24 (100)	14-511	96.5
<i>Hemibarbus labeo</i>	15	2 (13.3)	-	2.0
<i>Abbotina rivularis</i>	14	13 (92.9)	2-60	10.3
<i>Rhodeus ocellatus</i>	10	9 (90.0)	1-5	2.4
Subtotal	142	93 (65.5)	1-511	30.2
2019				
<i>Zacco platypus</i>	40	11 (27.5)	1-12	2.5
<i>Hemibarbus labeo</i>	34	14 (41.2)	1-10	2.2
<i>Pseudogobio esocinus</i>	28	10 (35.7)	1-10	2.6
<i>Hemibarbus longirostris</i>	8	6 (75.0)	1-8	3.0
<i>Pseudorasbora parva</i>	6	6 (100)	30-200	3.5
<i>Acanthorhodeus gracilis</i>	5	5 (100)	16-233	124.6
<i>Pungtungia herzi</i>	3	3 (100)	285-654	447.0
<i>Rhodeus ocellatus</i>	2	1 (50.0)	-	8.0
Subtotal	126	56 (44.4)	1-654	49.2
2020				
<i>Zacco platypus</i>	45	1 (2.2)	-	5.0
<i>Pseudorasbora parva</i>	26	16 (61.5)	1-290	65.3
<i>Hemibarbus labeo</i>	19	5 (26.3)	1-10	3.0
<i>Rhodeus ocellatus</i>	15	5 (33.3)	1-11	3.2
<i>Pungtungia herzi</i>	9	2 (22.2)	18-91	54.5
<i>Microphysogobio longidorsalis</i>	2	2 (100)	8-41	24.5
Subtotal	116	31 (26.7)	1-290	39.9
Total	384	180 (46.9)	1-654	37.8