

Multiplex PCR Detection of Waterborne Intestinal Protozoa: Microsporidia, *Cyclospora*, and *Cryptosporidium*

Seung-Hyun Lee¹, Migyo Joung², Sejoung Yoon², Kyoungjin Choi², Woo-Yoon Park³ and Jae-Ran Yu^{2,*}

¹Department of Microbiology, Konkuk University School of Medicine, Seoul 143-701, Korea; ²Department of Environmental and Tropical Medicine, Konkuk University School of Medicine, Seoul 143-701, Korea; ³Department of Radiation Oncology, College of Medicine, Chungbuk National University, Cheongju 361-763, Korea

Abstract: Recently, emerging waterborne protozoa, such as microsporidia, *Cyclospora*, and *Cryptosporidium*, have become a challenge to human health worldwide. Rapid, simple, and economical detection methods for these major waterborne protozoa in environmental and clinical samples are necessary to control infection and improve public health. In the present study, we developed a multiplex PCR test that is able to detect all these 3 major waterborne protozoa at the same time. Detection limits of the multiplex PCR method ranged from 10¹ to 10² oocysts or spores. The primers for microsporidia or *Cryptosporidium* used in this study can detect both *Enterocytozoon bieneusi* and *Encephalitozoon intestinalis*, or both *Cryptosporidium hominis* and *Cryptosporidium parvum*, respectively. Restriction enzyme digestion of PCR products with BsaBI or BsiEI makes it possible to distinguish the 2 species of microsporidia or *Cryptosporidium*, respectively. This simple, rapid, and cost-effective multiplex PCR method will be useful for detecting outbreaks or sporadic cases of waterborne protozoa infections.

Key words: *Cyclospora*, *Cryptosporidium*, microsporidia, multiplex PCR, waterborne protozoa

INTRODUCTION

Emerging waterborne protozoa, such as microsporidia, *Cyclospora*, and *Cryptosporidium*, have become a challenge to human health worldwide. These protozoans have several common characteristics biologically. Their major habitat is intestinal epithelial cells, and they are all intracellular parasites [1]. In addition, they produce infectious spores that are excreted from the hosts in their stools [2]. Although these protozoa are a concern for AIDS-infected individuals, they are gaining recognition as important infective organisms in immunocompetent individuals as well [1,3,4].

Of AIDS patients with chronic diarrhea, about 50% are diagnosed as infected with microsporidia, and 10-20% are infected with *Cryptosporidium parvum* [1]. A large waterborne outbreak of *C. parvum* occurred in 1993 [5], and outbreaks of *Cyclospora* that was due to consumption of imported raspberries have

been reported [6,7]. Microsporidia has been confirmed as a waterborne protozoon based on its detection in tertiary sewage effluent, surface water, and ground water [8]. Among the various genera in microsporidia, *Enterocytozoon bieneusi* and *Encephalitozoon intestinalis* are associated with gastrointestinal diseases in humans [8].

So far, there are very limited data available pertaining to the infection status of microsporidia, *Cyclospora*, and *Cryptosporidium* in Korea. Development of a simple, rapid, and economical detection method for these waterborne protozoa is urgently necessary to elucidate the infection rate in a community as well as individuals. Although previously Orlandi and Lampel [9] suggested PCR method for detection of these waterborne protozoa, their method has several disadvantages, such as demanding too much PCR reaction volume and providing no information on *Cryptosporidium hominis* infection.

We developed a multiplex PCR-based method, which is simple, rapid, and cost-effective for detecting these 3 kinds of major waterborne protozoa.

MATERIALS AND METHODS

E. intestinalis spores were purchased from ATCC (Cat. No. 50506; Manassas, Virginia, USA) and maintained through in

• Received 27 September 2010, revised 11 November 2010, accepted 18 November 2010.

* Corresponding author (maria205@kku.ac.kr)

© 2010, Korean Society for Parasitology

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

vitro culture using BS-C-1 cells (African green monkey kidney cells; Korean Cell Line Bank, Seoul, Korea) following the manufacturer's instructions. Briefly, Eagle's minimum essential medium (Sigma, St. Louis, Missouri, USA) with 2 mM L-glutamine, 1.5 g/L sodium bicarbonate, 0.1 mM non-essential amino acid, and 1.0 mM sodium pyruvate supplemented with 3% fetal bovine serum (GIBCO, Invitrogen, Carlsbad, California, USA) was used for maintenance of infection, and cells were incubated at 37°C with 5% CO₂. Oocysts of *Cryptosporidium parvum* (KKU isolate) were obtained from laboratory mice (C57BL6/J) that were infected and maintained as described [10]. The study was approved by the Animal Care and Use Committee of Konkuk University (Seoul, Republic of Korea).

For DNA extraction from fecal specimens, QIAquick stool mini kit (QIAGEN Inc., Valencia, California, USA) was used. For *E. intestinalis* and *C. parvum* DNA, each number of isolated spores or oocysts from 1×10^4 to 1×10^0 were spiked into 500 µg of uninfected human stool and mixed with 10 ml distilled water (DW). After washing with DW twice by centrifugation at 100 g, pellets were resuspended in ASL lysis buffer included in the kit. The samples were incubated at 70°C for 30 min. The procedure was executed in accordance with the manufacturer's recommendations. The extracted DNA was used as a template for PCR. DNA of *Cyclospora cayetanensis* was extracted from the stool of an infected patient who was the subject of a published case study in 2003 [11]. We cloned the 18S rRNA gene of *Cyclospora* using pGEM-T easy vector (Promega, Madison, Wisconsin, USA) and used it as a template DNA after diluted it equivalent to each number of parasite.

All of the PCR primers used in this study were designed using Primer-BLAST (<http://blast.ncbi.nlm.nih.gov>) and are shown in Table 1. The first-round multiplex PCR amplification was

performed in a 70-µl volume containing 5 µl of template DNA, 10 mM Tris-HCl (pH 8.3), 2 mM MgCl₂, 200 µM each of dATP, dCTP, dGTP, and dTTP, 0.7-1 µM of each primer (0.7 µM for microsporidia and 1 µM for *Cyclospora* and *Cryptosporidium*), and 5 U Taq DNA polymerase (Promega). The second-round multiplex PCR amplification was performed under the same conditions, except 2 µl of template DNA (product of the first-round PCR) was used in a 30-µl final volume. Both amplifications were performed using a C-1000 DNA thermal cycler (Bio-Rad, Hercules, California, USA) with initial denaturation at 94°C for 5 min, followed by 35 cycles at 94°C for 30 sec, 53°C for 30 sec, and 72°C for 90 sec and a final extension at 72°C for 10 min. The second-round PCR cycling conditions were identical to the first-round PCR except for the annealing temperature (55°C). Amplified DNA was analyzed by electrophoresis in a 2% (w/v) agarose gel stained with ethidium bromide (0.5 µg/ml) and visualized under ImageQuant300 (GE Healthcare, Giles, UK).

An aliquot (5 µl) of the second-round PCR product of *E. intestinalis* or *C. parvum* was used for enzyme digestion with BsaBI or BsiEI (New England BioLabs, Massachusetts, USA) at 60°C for 2 hr. DNA fragments were analyzed by electrophoresis in a 2% (w/v) agarose gel as described above.

RESULTS

From the first-round PCR amplification, we obtained the following products with the predicted sizes: 644-657 bp, 636 bp, and 415-427 bp for microsporidia, *C. cayetanensis*, and *Cryptosporidium*, respectively (data not shown). From the second-round PCR, the following predicted PCR products were amplified: 410-420 bp, 294 bp, and 171-183 bp for microsporidia,

Table 1. Primer sets used for multiplex PCR

| Parasite | PCR | Primer name | Sequence | PCR product size |
|--------------------------------|-----|-------------|------------------------------|------------------|
| Microsporidia | 1st | Mic A | 5'-GGAGCCTGAGAGATGGCT-3' | 644-657 bp |
| | | Mic E | 5'-AACGGCCATGCACCAC-3' | |
| | 2nd | Mic C | 5'-GGTGCCAGCAGCCGCGG-3' | 410-420 bp |
| | | Mic D | 5'-GCACAATCCACTCCT-3' | |
| <i>Cyclospora cayetanensis</i> | 1st | CYCF1E | 5'-TACCCAATGAAAACAGTTT-3' | 636 bp |
| | | CYCR2B | 5'-CAGGAGAAGCCAAGGTAGG-3' | |
| | 2nd | CYCF3E | 5'-CCTTCCGCGCTTCGCTGCGT-3' | 294 bp |
| | | CYCR4B | 5'-CGTCTTCAAACCCCTACTG-3' | |
| <i>Cryptosporidium</i> spp. | 1st | cp2415F | 5'-CCCACGCGAAGTTGAAGTAAC-3' | 415-427 bp |
| | | cp2415R | 5'-CTTAGTTGCTTGCTTGAGTTGG-3' | |
| | 2nd | cp2171F | 5'-CAACCCAGAAGTTGAGGTT-3' | 171-183 bp |
| | | cp2171R | 5'-CTAGTATGCTTCAGACCATGAG-3' | |

C. cayetanensis, and *Cryptosporidium*, respectively (Fig. 1). The lower detection limit of nested PCR amplification that was performed with template DNA from each individual protozoa was 10^2 spores for microsporidia, 10^0 oocysts for *Cyclospora*, and 10^1 oocysts for *Cryptosporidium* (Fig. 1). The lower detection limit of multiplex PCR amplification that was performed with mixed DNA from each protozoan was 10^2 for microsporidia and *Cyclospora* and 10^1 for *Cryptosporidium* (Fig. 2).

The primers for microsporidia and *Cryptosporidium* used in the present study are capable of detecting both *E. bienersi* and *E. intestinalis*, and *C. parvum* and *C. hominis*, respectively. Restriction enzyme digestion of the resulting nested PCR products can distinguish the 2 kinds of microsporidia and *Cryptosporidium* spp. Restriction enzyme digestion with BsaBI showed 2 frag-




Fig. 1. Sensitivity of nested PCR primer sets for microsporidia (A), *Cyclospora cayetanensis* (B), and *Cryptosporidium* (C). M, DNA marker; N, negative control (DW).

mented bands of 167 and 253 bp in *E. intestinalis* (Fig. 3A), and with BsiEI showed no fragmented bands in *C. parvum* (Fig. 3B). Whereas it was confirmed with Clone Manager 6 (Sci-Ed, North Carolina, USA) that there will be no fragmented bands in *E. bienersi* with BsaBI digestion and 2 fragmented bands of 48 and 137 bp in *C. hominis* with BsiEI digestion. Each primer set for microsporidia, *C. cayetanensis*, and *Cryptosporidium* used in the present study showed no cross-reactivity with each of the other parasite DNA (Fig. 4).

DISCUSSION

There are various detection methods currently available for microsporidia, *Cyclospora*, and *Cryptosporidium*. These include




Fig. 2. Sensitivity of multiplex nested PCR for microsporidia, *Cyclospora cayetanensis*, and *Cryptosporidium*. Serially diluted template DNA from each type of protozoan was mixed, and 3 kinds of primer sets for each type were included in 1 reaction tube. M, DNA marker; N, negative control (DW).




Fig. 3. Restriction enzyme digestion of nested PCR products of microsporidia digested with BsaBI (A) and *Cryptosporidium parvum* digested with BsiEI (B). M, DNA marker; Ei, *E. intestinalis*; Cp, *C. parvum*.




Fig. 4. Specificity of multiplex nested PCR primers. No cross-reacted PCR bands were detected when the 3 primer sets were paired with DNA from the different protozoa. Lane 4, negative control (DW).

morphological examination after specific staining, fluorescence microscopy after specific antibody labeling, and molecular diagnosis with such methods as PCR [2,12,13]. Although it is inexpensive and fast, the classical staining method depends on highly trained experts for accurate diagnosis. Immunofluorescence-based antibody-labeling methods depend on the sensitivity and specificity of antibodies against parasites for the accurate diagnosis, and cross-reactivity or nonspecific labeling can be a challenge. Multiplex PCR would eliminate the need for a highly trained expert and would reduce the diagnostic time by being able to catch 3 kinds of protozoa at the same time. If well-designed multiplex PCR having a high sensitivity and specificity is available, it would be an excellent diagnostic method for these waterborne protozoa.

The primers used for *Cyclospora* in this study were designed by Relman et al. [14]. Orlandi and Lampel [9] reported a sensitivity level for *Cyclospora* that was as low as 3 oocysts using nested PCR with the same primer set, which is similar to the level of sensitivity in our study. In addition, Orlandi and Lampel [9] described a detection sensitivity of *C. parvum* and *E. intestinalis* of as few as 10 oocysts from singleplex PCRs with each parasite DNA [9]. Their detection sensitivity for *E. intestinalis* is 10-fold higher than our results; however, they did not report the sensitivity of multiplex PCR using mixed DNA of each protozoan. Furthermore, agarose gel electrophoresis analysis of PCR products from both the first and second rounds of PCR is necessary to confirm the detection results with the method described by Orlandi and Lampel [9], because that method amplified the *C.*

parvum target only during the first round of PCR.

Generally the sensitivity of multiplex PCR is not exactly the same with individual PCR done with each parasite DNA. Multiplex PCR developed in the present study showed the same detection sensitivity with that of individual PCR in case of microsporidia and *Cryptosporidium*. Agarose gel electrophoresis of the second-round PCR products could provide all of the detection results, when the multiplex PCR tests reported here were applied.

Another advantage of the multiplex PCR test developed in the present study is that the 2 species of microsporidia, such as *E. bienersi* and *E. intestinalis*, and the 2 species of *Cryptosporidium*, such as *C. parvum* and *C. hominis*, can be differentiated after PCR by restriction enzyme digestion. We did not show restriction enzyme-digested PCR products of *E. bienersi* and *C. hominis* in the present study as we did not secure DNAs of these 2 organisms. Instead, we confirmed the results of restriction enzyme digestion of these 2 organisms after PCR using a restriction enzyme cutting software program (Clone Manager 6).

Although there is a well-established genomic database for *Cryptosporidium* (<http://cryptodb.org>) that was established through the efforts of many devoted and excellent scientists, there are still only limited genomic data available for *Cyclospora* and microsporidia. Therefore it is quite difficult to develop specific primers for these 2 protozoa. The difficulty is especially challenging for *Cyclospora* because PCR detection of the 18S rRNA gene could be confused with that of *Eimeria* spp., which show 98% similarity with the 18S rRNA gene sequence of *Cyclospora*. Thus sequence analysis after multiplex PCR should be performed for the accurate differential diagnosis of *Cyclospora* and *Eimeria* spp.

In conclusion, the multiplex PCR test developed in the present study can detect microsporidia, *Cyclospora*, and *Cryptosporidium*, all of which cause severe waterborne diarrheal disease. The test is very simple and rapid and offers high sensitivity and specificity. This test could help improve detection of diarrheal outbreaks or sporadic diarrheal disease that is due to infection with these major waterborne protozoa which have been considered as unknown etiology for the difficulty in detection previously.

ACKNOWLEDGEMENTS

This research was supported by the Basic Science Research Program through the National Research Foundation of Korea (RF) funded by the Ministry of Education, Science, and Tech-

nology (2010-0017321).

REFERENCES

1. Goodgame RW. Understanding intestinal spore-forming protozoa: cryptosporidia, microsporidia, isospora, and cyclospora. *Ann Intern Med* 1996; 124: 429-441.
2. O'Donoghue PJ. *Cryptosporidium* and cryptosporidiosis in man and animals. *Int J Parasitol* 1995; 25: 139-195.
3. Sandfort J, Hannemann A, Gelderblom H, Stark K, Owen RL, Ruf B. *Enterocytozoon bieneusi* infection in an immunocompetent patient who had acute diarrhea and who was not infected with the human immunodeficiency virus. *Clin Infect Dis* 1994; 19: 514-516.
4. Lee JK, Song HJ, Yu JR. Prevalence of diarrhea caused by *Cryptosporidium parvum* in non-HIV patients in Jeollanam-do, Korea. *Korean J Parasitol* 2005; 43: 111-114.
5. Mac Kenzie WR, Hoxie NJ, Proctor ME, Gradus MS, Blair KA, Peterson DE, Kazmierczak JJ, Addiss DG, Fox KR, Rose JB, et al. A massive outbreak in Milwaukee of *Cryptosporidium* infection transmitted through the public water supply. *N Engl J Med* 1994; 331: 161-167.
6. Herwaldt BL, Ackers ML. An outbreak in 1996 of cyclosporiasis associated with imported raspberries. The *Cyclospora* Working Group. *N Engl J Med* 1997; 336: 1548-1556.
7. Ortega YR, Roxas CR, Gilman RH, Miller NJ, Cabrera L, Taquiri C, Sterling CR. Isolation of *Cryptosporidium parvum* and *Cyclospora cayetanensis* from vegetables collected in markets of an endemic region in Peru. *Am J Trop Med Hyg* 1997; 57: 683-686.
8. Dowd SE, Gerba CP, Pepper IL. Confirmation of the human-pathogenic microsporidia *Enterocytozoon bieneusi*, *Encephalitozoon intestinalis*, and *Vittaforma corneae* in water. *Appl Environ Microbiol* 1998; 64: 3332-3335.
9. Orlandi PA, Lampel KA. Extraction-free, filter-based template preparation for rapid and sensitive PCR detection of pathogenic parasitic protozoa. *J Clin Microbiol* 2000; 38: 2271-2277.
10. Lee SU, Joung M, Ahn MH, Huh S, Song H, Park WY, Yu JR. CP2 gene as a useful viability marker for *Cryptosporidium parvum*. *Parasitol Res* 2008; 102: 381-387.
11. Yu JR, Sohn WM. A case of human cyclosporiasis causing traveler's diarrhea after visiting Indonesia. *J Korean Med Sci* 2003; 18: 738-741.
12. Didier ES. Microsporidiosis: an emerging and opportunistic infection in humans and animals. *Acta Trop* 2005; 94: 61-76.
13. Shields JM, Olson BH. *Cyclospora cayetanensis*: a review of an emerging parasitic coccidian. *Int J Parasitol* 2003; 33: 371-391.
14. Relman DA, Schmidt TM, Gajadhar A, Sogin M, Cross J, Yoder K, Sethabutr O, Echeverria P. Molecular phylogenetic analysis of *Cyclospora*, the human intestinal pathogen, suggests that it is closely related to *Eimeria* species. *J Infect Dis* 1996; 173: 440-445.

